



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b>  <b>A61K 31/70</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 93/25209</b>  <b>(43) International Publication Date:</b> 23 December 1993 (23.12.93)
<b>(21) International Application Number:</b> PCT/HU93/00038 <b>(22) International Filing Date:</b> 17 June 1993 (17.06.93)  <b>(30) Priority data:</b> P 92 02040 18 June 1992 (18.06.92) HU  <b>(71) Applicant (for all designated States except US):</b> FABULON RG KOZMETIKAI KFT. [HU/HU]; Gyömrői út 12, H- 1103 Budapest (HU).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> HANGAY, György [HU/HU]; Rajk László u. 35-37, H-Budapest XIII (HU). KELEN, András [HU/HU]; Somfa köz 6, H-1106 Budapest (HU). KESERŰ, Péter [HU/HU]; Párkány u. 20, H-1138 Budapest (HU). DEMETER, Vodnár, János [HU/HU]; Rudnay Sándor tér 6, H-2500 Esztergom (HU). FRIEDMAN, Éva [HU/HU]; Fehérvári u. 147, H-1119 Budapest (HU). ALBERTI, Éva [HU/HU]; Hild József u. 2, H-2500 Esztergom (HU). KOVÁCS, György [HU/HU]; József u. 27, H-Budapest XVIII (HU). PUTZ, Pál [HU/HU]; Kun Béla ltp. 8/3, H-2500 Eszter- gom (HU). KARDICS, Izabella [HU/HU]; Bánomi ltp. 6, H-2500 Esztergom (HU).		<b>(72) Inventor:</b> MÉSZÁROS, Józsefné (deceased).  <b>(74) Agent:</b> DANUBIA; Bajcsy Zsilinszky út 16, H-1051 Bu- dapest (HU).  <b>(81) Designated States:</b> AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, JP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> COMPOSITIONS FOR THE TREATMENT AND PROPHYLAXIS OF INFLAMMATIONS AND DERMA- TOSSES INDUCED BY VIRUSES AND PROCESS FOR THE PREPARATION THEREOF  <b>(57) Abstract</b>  The invention relates to new cosmetic and/or medicinal compositions which beneficially affect virus-induced infections and inflammations of the skin and mucosa and promote the regeneration of the dermatoses caused, i.e. the compositions are able to restore the initial conditions of the skin.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LJ	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TC	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

COMPOSITIONS FOR THE TREATMENT AND PROPHYLAXIS OF  
INFLAMMATIONS AND DERMATOSES INDUCED BY VIRUSES AND  
PROCESS FOR THE PREPARATION THEREOF

5           The invention relates to new cosmetic and/or  
medicinal compositions which beneficially affect  
virus-induced infections and inflammations of the  
skin and mucosa and promote the regeneration of the  
dermatoses caused, i.e. the compositions are able to  
10 restore the initial conditions of the skin.

The compositions of the invention are special-  
ly suitable for the treatment of infections and  
dermatoses induced by herpes virus.

Several patents report on products suitable  
15 for the treatment of herpes. The US patent  
specification No. 317976 recommends the hydrophilic  
ointment tricontanol. Tricontanol (melissyl alcohol,  
myricyl alcohol,  $\text{CH}_3(\text{CH}_2)_{28}\text{CH}_2\text{OH}$ ) alleviates pain,  
heals the wounds developing in acne and perioral  
20 dermatitis, exerts antiinflammatory action and  
restores the normal lipid content of the skin. In  
the Swiss patent specification No. 005810 (CIBA) an  
oral stick formulation containing heparin and  
dissociating zinc salt are proposed. The Israeli  
25 patent specification No. 065184 (see British patent  
specification No. 029749 and US patent specification  
No. 4,700,293, too) recommends a combination of  
glycyrrhizin and steroids (e. g. triamcinolon) or  
viricidal agents (e. g. idoxuridine). The first is  
30 proposed for the treatment of aphtha with adjuvant  
stomatitis while the latter one for oral, nasal or  
genital herpes. The Roumanian patent specification  
No. 098473 describes a mixture of poplar bud  
extract, arnica blossom tincture and melilot oil.  
35 The Hungarian patent specification No. 001224

- 2 -

reports on antiherpetic 2'-chloro-2'-deoxyuridines substituted in position 5.

The cited patents also confirm that in this field of indication the use of natural substances is becoming popular world-wide, too. The present invention is also a representative of this trend of development, and is based on the observation that active ingredient concentrates can be prepared from particular natural substances by a specific process which can be used, alone or in combination, for the production of cosmetic agents or medicinal cosmetics of improved activity. Their activity can be further increased by applying adjuvants or specific formulations.

The invention relates to compositions of manifold application with viricide, antiinflammatory and regenerating activity wherein the concentrations of the ingredients given % by weight are as follows:

20	Glycyrrhizin, preferably in the form of sweet-root extract	0.005-0.4
	Rosemary acid, preferably in the form of plant extract	0.005-0.4
	Thyme oil (thymol content min. 20 %)	0.001-0.5
25	Allantoin	0-0.5
	Polydocanol	0-5.0
	$\alpha$ -Bisabolol	0.005-0.5
	Soluble collagen	0.001-0.05
	Irgasan OP 300	0-0.5
30	Ethanol	0-90.

The compositions may contain further carriers, additives and auxiliary materials usually applied in the preparation of cosmetic or medicinal compositions, preferably water, propyleneglycol, acrylamide-sodium acrylate copolymer, PEG-7,

- 3 -

glycerol cocoate and isooctyl nonanoate, as well as solubilizing agents, lipids, preservatives and fragrance components in hydrogel, lotion-ointment, liquid emulsion, spray, stick or drop formulation.

5       As regards the constituents of the composition it is known from literature sources that rosemary acid (caffeoyl- $\alpha$ -hydroxy-dihydrocaffeic acid), together with quinic acid, chlorogenic acid and caffeic acid, is present in plants of the labiate family (Lamiaceae), mainly in *Melissa officinalis*, *Hyssopus officinalis*, *Rosmarinus officinalis* and *Salvia officinalis*. Its virostatic, antibacterial and antimycotic potency is reported by I. K. Heimann and W. Schultze in the *Zeitschrift für Phytotherapie* Sept. 1988, 82-83.

15       Because of their sweet taste the sweet root (*Glycyrrhiza glabra*) extracts are used as sweeteners to improve taste and as expectorants (B. Issekutz: (Drug prescription, *Gyógyszerrendelés* p. 600 *Medicina*, Budapest, 1979). The regular use of standard preparations may reduce the increased tension of the stomach wall and the inflammation of gastric mucosa (Gizella Verzár Petri: *Gyógynövények a gyógyászatban* (Medicinal plants in therapy), p. 92, *Medicina*, Budapest, 1984). Soviet authors [I. A. Murajev 25 et al.: *Farmacia* (Moscow), 94 (2), 84, 25 1979] found it to be more or less equiactive with prednisolone in contact allergy and dermatitis. Glycyrrhizin, the major constituent of sweet root has mineralocorticoid, antiinflammatory, antiedema and epithelizing (wound healing) properties.

30       The volatile oil of thyme (*Thymus vulgaris*) contains mainly thymol, carvacrol, p-cymol, furthermore linalool, borneol, geraniol and  $\beta$ -pinene. Due to its thymol content the thyme oil has viricide effect and inhibits a major part of wound bacteria even at 3000fold dilution (Gizella Verzár

- 4 -

Petri: Medicinal plants in therapy, Medicina, Budapest, 1984, p. 74.).

The  $\alpha$ -bisabolol is one of the major components of camomile (*Matricaria chamomilla*). Primarily it  
5 has antiinflammatory effect and promotes granulation.

The allantoin promotes the skin-regeneration and hydration.

The collagens, which can be found in the skin, also play role in hydration, but based on other  
10 mechanism.

The polydocanol is ingredient of numerous medicinal and cosmetic composition. It has analgetic and local anaesthetic effect.

According to the literature individual compo-  
15 nents of the preparation of the present invention are applied, though with varying frequency, in medical practice, but no preparation which has similar composition as that of the invention and is produced with a process similar to the disclosed  
20 one. The product(s) are highly potent, have a broad spectrum of activity and promote the rapid and full regeneration of the affected skin areas.

The preparation of individual medicinal plant extracts, used in the preparation, as well as the  
25 composition of the compositions are summed up as follows:

Characterization of components:

1.) Dry extract with rosemary acid content:

An aqueous alcohol (methanol, ethanol, iso-  
30 propanol) extract is prepared from the leaves of a plant from Laminaceae family, e.g. rosemary, melilot (*Melissa officinalis*), *Salvia officinalis* and *Hyssopus officinalis*, etc. After removing the solvent the residue is acidified to pH 2-3 with an  
35 inorganic acid, the mixture is extracted with a

- 5 -

water immiscible solvent (ether, chloroform, dichloromethane, ethyl acetate, etc.), the layers are separated and the solvent is removed from the organic layer yielding a dry extract.

5

2.) Dry extract containing glycyrrhizin:

The sweet root is extracted with water or aqueous alcohol (methanol, ethanol, isopropanol) and the solvent is removed. The aqueous solution is  
10 acidified to pH 2-3 with an inorganic acid, the precipitated mixture is filtered, the precipitate is dried, dissolved in ethanol, the pH of the solution is adjusted to 7-8 with ammonia, filtered and evaporated to dryness.

15 3.) Thyme oil. *Aetheroleum thymi* (Hungarian Pharmacopoea No. 7.)

Purified volatile oil prepared from the blossoming, fresh branch tops of the garden thyme (*Thymus vulgaris*) with steam distillation. The  
20 product has a phenol content not less than 20.0 % (v/w) expressed in thymol.

The examination of the effect of the compositions:

The product is specifically beneficial for symptoms  
25 of the mucosa (oral and genital) induced by herpes simplex virus. Applied in the early stage the development of erythema and edema is suppressed and recovery is accelerated. In recurrent cases, applied before the development of symptoms, superinfection is fully inhibited.  
30 bited. Furthermore, subjective symptoms, such as itching and stretching, are alleviated, and recovery is rapid and traceless. The skin becomes refreshed and hydrated.

Two figures are shown to demonstrate the efficacy of the product. Figure 1 presents the results of the comparative study of the preparation of the invention (herpes  
35

- 6 -

gel prepared according to procedure 1) and that of the commercial control products, together with the duration of recovery of untreated cases. The following control preparations were tested in the study (Figure 1 shows the mean values of the results obtained):

1.) Medicinal gel and paint formulation containing the extract of 15 % (w/w) *Chelidonium herba*, 3 % (w/w) *Menthae piperita folium*, 1 % (w/w) *Calendula officinalis*, 2 % (w/w) *Thymi herba* and 2 % (w/w) *Millefolii herba*, prepared with 50 % (w/w) ethanol.

2.) Medicinal suspension ointment formulation containing 0.8 % (w/w) of 3-(2-deoxy-D-ribofuranosyl)-5-isopropyluracil.

3.) Medicinal alcoholic gel formulation containing 0.2 % (w/w) *primicium sulfuricum* and 2 % (w/w) *lidocain*.

4.) Painting agent containing 0.5 % (w/w) of zinc sulfate widely used in magistral preparations.

The compositions according to the invention reduced the duration of symptom free recovery by 3-4 days compared to the control products and by 7-9 days compared to the untreated control cases. Figure 2 presents the distribution of major recovery categories in the case of the preparation of the invention. According to the chart in about 30 % of the subjects crusts or blisters failed to develop and symptoms were eliminated within 2 days. In about 60 % of the patients only a thin and flexible crust was formed and wound healing was accelerated. Finally in about 10 % of the cases the rate of recovery, both as regards symptoms and wound healing, was similar to that observed with the control preparations. Further observations confirmed that in the case of herpes simplex infections the



product prolonged the interval elapsing till the recurrence of the disease. The symptom free periods may even increase from one to two weeks to 2 months.

From the tests performed the diagram of the hydrating effect of the gel containing the entire active ingredient combination, taken with a SKIN Resistance Meter, is shown in Figure 3.

Preferred formulations of the preparations according to the invention are listed as follows:

In the major part of indications the most advantageous formulation proved to be the rapidly absorbing and penetrating gel form which is easy to handle. The formulations specified (1-3) are partly usual gels and partly microemulsions [values in the table are expressed in % by weight (w/w)]:

<u>Herpes gel formulations</u>		<u>1</u>	<u>2</u>	<u>3</u>
Acrylic acid/Na acrylate				
copolymer (Hostacerin PN 73)		2.0	1.5	
POE(45) hydrogenated ricinol				
acid triglyceride				
(Cremophor RH 40)		2.0	1.5	
POE(7) glycerol cocoate				
(Cetiol HE)		2.0	5.0	
Isooctadecyl isononanoate				
(PCL liquid)		2.0	0.3	
Propyleneglycol		8.0	10.0	12.0
Ethanol		30.0	30.0	16.0
POE(9) lauryl alcohol				
(Polidokanol)		2.0	1.0	2.0
Polyoxyethylene/polyoxypropylene block polymer				
(Lutrol FC 127)				19.0
Allantoin		0.1	0.2	
$\alpha$ -Bisabolol		0.1	0.05	0.1
1 w/w % Collagen solution		0.5	1.0	0.5

- 8 -

	Triclosan (Irgasan OP 300)		0.1	0.1	
	Glycyrrhizin		0.1	0.2	0.15
	Rosemary acid		0.1	0.2	0.15
	Thyme oil		0.02	0.025	0.02
5	Distilled water	ad	100	100	100

The gels are prepared by dissolving allantoin in the bulk of water, then in some cases (formulations 1-2) also the copolymer, using slow stirring.

Polydocanol, and in some cases (formulations 1-2) also Irgasan, are dissolved in part of the alcohol. The remaining volumes of alcohol and water are combined and the formerly solubilized thyme oil is added to the mixture. The phases are stirred at low r. p. m., then the lipid additives are poured to the mixture under stirring in the following order: solution of rosemary acid and glycyrrhizin in propyleneglycol and solution of  $\alpha$ -bisabolol and collagen. After the addition stirring is continued for further 5-10 minutes.

Ointment is also a formulation of choice, see formulations 4 to 6 [values in the table are expressed in % by weight (w/w)]:

25	<u>Herpes ointment formulations</u>	<u>4</u>	<u>5</u>	<u>6</u>
	Liquid paraffin	18.0	18.0	20.0
	Stearin	7.0	8.0	9.0
	Polyoxyethylene fatty acid ester	6.0	7.0	8.0
30	Isooctyl stearate	4.0	3.0	3.0
	Propyleneglycol	4.0	7.5	12.0
	POE-9 lauryl alcohol (polydocanol)		1.0	2.0
	Allantoin	0.1	0.1	0.2
35	$\alpha$ -Bisabolol	0.1	0.15	0.05

- 9 -

	1 w/w % Collagen solution	0.5	0.5	1.0
	Triclosan (Irgasan DP 300)		0.1	0.1
	Glycyrrhizin	0.05	0.1	0.2
	Rosemary acid	0.05	0.1	0.2
5	Thyme oil	0.03	0.04	0.05
	Distilled water	ad	100	100

Ointments are prepared by melting the lipid components and the emulsifier - liquid paraffin, stearin, isooctyl stearate or polyoxyethylene (POE) fatty acid ester, resp. - and heating the mixture to about 70°C. Thereafter the polydocanol is added, and, if desired (formulations 5 and 6), Irgasan. Allantoin is dissolved in water then the aqueous phase is heated to about 70°C.

15 After completed homogenization, under steady cooling and moderate stirring, at about 60°C the solutions of rosemary acid and glycyrrhizin in propyleneglycol, solutions of  $\alpha$ -bisabolol and collagen, then at temperatures lower than 50°C thyme oil are added. Stirring is continued till the mixture becomes cool (about 30°C).

Alcoholic paint, drop or spray are also formulations of choice. The composition of formulations 7 to 9 are listed in the following.

25

<u>Herpes formulations</u>		<u>7</u>	<u>8</u>	<u>9</u>
	Ethanol	20.0	30.0	
	Propylene glycol	5.0	8.0	40.0
	POE(45) hydrogenated ricinol			
30	acid triglyceride			
	(Cremophor RH 40)	1.0	2.0	3.0
	Triclosan (Irgasan OP 300)	0.05	0.1	
	Allantoin	0.05	0.1	0.2
	$\alpha$ -Bisabolol	0.05	0.2	0.1
35	1 w/w % Collagen solution	1.0	0.5	0.5

- 10 -

Glycyrrhizin		0.05	0.1	0.2
Rosemary acid		0.05	0.1	0.2
Thyme oil		0.02	0.05	0.05
Distilled water	ad	100	100	100

5

The solutions are prepared by solubilizing first the thyme oil, preferably with POE(45) hydrogenated ricinol acid triglyceride. Irgasan is dissolved in 10 alcohol, and allantoin in the bulk of water. The phases prepared in this way are mixed, then, under constant stirring the solution of  $\alpha$ -bisabolol and collagen (formulations 7 and 8) is added to the system, finally the propyleneglycol solution of 15 glycyrrhizin and rosemary acid. Stirring is continued for a couple of minutes after the addition.

Alcoholic painting composition and aerosol foam or spray composition using an appropriate carrier gas can be also prepared.

What we claim is

5

1. Compositions for the treatment and prophylaxis of inflammations and dermatoses of the skin induced by viruses characterized in that containing ingredients given in % by weight as

10 follows:

	Glycyrrhizin	0.005-0.4
	Rosemary acid	0.005-0.4
	Thyme oil (thymol content min. 20 %)	0.001-0.5
	Allantoin	0-0.5
15	Polydocanol	0-5.0
	$\alpha$ -Bisabolol	0.005-0.5
	Soluble collagen	0.001-0.05
	Irgasan OP 300	0-0.5
	Ethanol	0-90

20 furthermore auxiliary materials usually applied in the preparation of cosmetic and medicinal compositions.

2. Compositions as claimed in claim 1 characterized in that containing as auxiliary materials water, propyleneglycol, acrylamide/sodium  
25 acrylate copolymer, PEG-7-glyceryl cocoate, isooctyl nonanoate, solubilizing agents, lipids and preservatives.

3. Compositions as claimed in either of claims 1 or 2 for the treatment and prophylaxis of viral infections and inflammations characterized in that the composition  
30 is formulated as a liquid, drop, aerosol, foam, ointment, emulsion or gel formulation.

4. A process for preparing compositions suitable for the treatment and prophylaxis of inflammations and dermatoses of the skin induced by viruses which comprises  
35 mixing the ingredients given in % by weight as follows:

	Glycyrrhizin	0.005-0.4
	Rosemary acid	0.005-0.4
	Thyme oil (thymol content min. 20 %)	0.001-0.5
5	Allantoin	0-0.5
	Polydocanol	0-5.0
	$\alpha$ -Bisabolol	0.005-0.5
	Soluble collagen	0.001-0.05
	Irgasan OP 300	0-0.5
10	Ethanol	0-90

furthermore auxiliary materials usually applied in the preparation of cosmetic and medicinal composition and converting them to cosmetic or medicinal compositions.

5. A process as claimed in claim 4 which  
 15 comprises using water, propyleneglycol, acrylamide sodium acrylate copolymer, PEG-7-glyceryl cocoate, isooctyl nonanoate furthermore solubilizing agents, lipids and preservatives as auxiliary materials.

6. A process as claimed in either of the claims  
 20 4 or 5 which comprises converting the compositions to liquid, drop, aerosol, foam, ointment, emulsion or gel formulations.

7. Therapeutic process for the treatment and  
 prophylaxis of infections, inflammations and  
 25 dermatoses of the skin and mucosa induced by viruses which comprises treating the infected body surface or the surface susceptible to infection with an effective dose of the preparation containing ingredients given in % by weight as follows:

30	Glycyrrhizin	0.005-0.4
	Rosemary acid	0.005-0.4
	Thyme oil (thymol content min. 20 %)	0.001-0.5
	Allantoin	0-0.5
	Polydocanol	0-5.0
35	$\alpha$ -Bisabolol	0.005-0.5

Soluble collagen

0.001-0.05

Irgasan OP 300

0-0.5

Ethanol

0-90

furthermore auxiliary materials usually applied in the  
5 preparation of cosmetic and medicinal composition.

1/3




Fig.1



2/3




Fig. 2




Fig. 3

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/HU 93/00038

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Patent Abstracts of Japan, unexamined applications Section C, Volume 16, no. 20, issued 1992, January 20, The Patent Office Japanese Government, page 17 C 902, the abstract no. 03-236 323 (KOBAYASHI KOSE CO LTD)..	1
A	Patent Abstracts of Japan, unexamined applications, Section C, Volume 16, no. 207, issued 1992, May 18, The Patent Office Japanese Government, page 23 C 941, the abstract no. 04-36 215 (SUNSTAR INC).	1
A	FR, A1, 2 614 787 (POLA KASEI KOGYO KABUSHIKI KAISHA) 10 November 1988 (10.11.88), claims 1,2,5. -----	1,5

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/HU 93/00038

## A. CLASSIFICATION OF SUBJECT MATTER

IPC<sup>5</sup>: A 61 K 31/70

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC<sup>5</sup>: A 61 K 31/70

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS, WPIL; glycyrrhizin?  
glycyrrhizinic w acid ) skin

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB, A, 2 167 296 (YISSUM RESEARCH AND DEVELOPEMENT COMPANY OF THE HEWBREW UNIVERSITY OF JERUSAMLEM) 29 May 1986 (29.05.86), pages 3,4, examples 8-12.	1,7
A	EP, A1, 0 487 404 (KABUSHIKI KAISHA HAYASHIBARA SEIBUTSU KAGAKU KENKYUJO) 27 May 1992 (27.05.92), claim 6.	1
A	GB, A, 2 071 494 (KANEBO LTD. & MARUZEN KASEI CO. LTD.) 23 September 1981 (23.09.81), abstract.	1
A	Patent Abstracts of Japan, unexamined applications, Section C, Volume 8, no. 98, issued 1984, May 09, The Patent Office Japanese Government, page 101 C 221, the abstract no. 59-13 716 (RAION K.K.).	1
A	Patent Abstracts of Japan, unexamined applications, Section C, Volume 12, no. 467, issued 1988, December 07, The Patent Office Japanese Government, page 120 C 550, the abstract no. 63-188 628 (SHISEIDO CO.)	1

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date  
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

26 August 1993 (26.08.93)

Date of mailing of the international search report

07 September 1993 (07.09.93)

Name and mailing address of the ISA/ AT  
AUSTRIAN PATENT OFFICE  
Kohlmarkt 8-10  
A-1014 Vienna  
Facsimile No. 0222/53424/535

Authorized officer

Schnass e.h.

Telephone No. 0222/5337058/35

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International application No.

PCT/HU 93/00038

Im Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
GB A 2167296		GB A0 8429749 GB A1 2167296 GB B2 2167296	03-01-85 29-05-86 16-11-88
EP A1 487404	27-05-92	AU A1 87975/91 JP A2 4182413 JP A2 4182412	21-05-92 30-06-92 30-06-92
GB A 2071494		FR A1 2477413 FR B1 2477413 GB A1 2071494 GB B2 2071494 JP A2 56125306 JP B4 63001921 US A 4481187 JP A2 57062215 JP B4 63020406	11-09-81 22-02-85 23-09-81 31-05-84 01-10-81 14-01-88 06-11-84 15-04-82 27-04-88
FR A1 2614787	10-11-88	JP A2 63275506	14-11-88